arXiv:1301.6332v1 [math.RA] 27 Jan 2013

Integer-valued polynomials over matrices and divided differences

Giulio Peruginelli[*]

January 29, 2013

Abstract

Let D be an integrally closed domain with quotient field K and n a positive integer. We give a characterization of the polynomials in $K[X]$ which are integer-valued over the set of matrices $M_n(D)$ in terms of their divided differences. A necessary and sufficient condition on $f \in K[X]$ to be integer-valued over $M_n(D)$ is that, for each k less than n, the k-th divided difference of f is integral-valued on every subset of the roots of any monic polynomial over D of degree n. If in addition D has zero Jacobson radical then it is sufficient to check the above conditions on subsets of the roots of monic irreducible polynomials of degree n, that is, conjugate integral elements of degree n over D.

Keyword Integer-valued polynomial, Divided differences, Matrix, Integral element, Polynomial closure. MSC Classification codes: 13B25, 13F20, 11C20.

1 Introduction

Let D be a domain with quotient field K. We denote by $M_n(D)$ the D-algebra of $n \times n$ matrices with entries in D. In this paper we are interested in the ring of polynomials in $K[X]$ which are integer-valued over $M_n(D)$ (see [5], for a general reference), namely:

$$\mathrm{Int}(M_n(D)) \doteqdot \{f \in K[X] \mid f(M) \in M_n(D), \forall M \in M_n(D)\}.$$

In [3] the authors consider the overring of integer-valued polynomials which are integer-valued over the subset $T_n(D) \subset M_n(D)$ of triangular matrices over D, namely the ring $\mathrm{Int}(T_n(D)) \doteqdot \{f \in K[X] \mid f(T) \in M_n(D), \forall T \in T_n(D)\}$. They characterize a polynomial $f(X)$ of this ring in terms of its divided differences, proving that up to the order $n-1$ the k-th divided difference $\Phi^k(f) \in K[X_1, \ldots, X_{k+1}]$ is integer-valued over D^{k+1}, that is $\Phi^k(f)(D^{k+1}) \subset D$. See section 3 of this paper for the definition of divided difference of a polynomial.

We give here an analogous characterization of the polynomials in $\mathrm{Int}(M_n(D))$. It turns out that, for every $0 \leq k < n$, $\Phi^k(f)$ is integral-valued on every subset of $k+1$ elements of the roots

---

[*] Institut für Analysis und Comput. Number Theory, Technische Univ., Steyrergasse 30, A-8010 Graz, Austria. E-mail: peruginelli@math.tugraz.at.

of any monic polynomial $p \in D[X]$ of degree n. More precisely, given such a polynomial $p(X)$, let $\alpha_1, \ldots, \alpha_n$ be its roots in a fixed algebraic closure of K. Then $\Phi^k(f)(\alpha_1, \ldots, \alpha_{k+1})$ is integral over D, for every possible choice of the subset $\{\alpha_i\}_{i=1}^{k+1}$ of the roots of $p(X)$. This property of $\Phi^k(f)$ holds for every polynomial $p(X)$ chosen as before. Notice that the α_i's are integral over D since $p(X)$ is monic (they are not conjugate over D, if $p(X)$ is reducible). These properties about the divided differences of $f(X)$ are sufficient and necessary for $f(X)$ to belong to $\mathrm{Int}(M_n(D))$. This characterization generalizes the previous one about integer-valued polynomials over triangular matrices. In fact, for such a polynomial $f(X)$ and for all the relevant k's, $\Phi^k(f)$ is integer-valued on every subset of $k+1$ roots of every monic polynomial over D of degree n which is totally split over D.

We give some preliminary notation for the next sections. Let $\overline{K}$ be a fixed algebraic closure of K and let $\overline{D}$ be the integral closure of D in $\overline{K}$. Given $M \in M_n(D)$, we denote by $p_M(X)$ the characteristic polynomial of M and by $N_{D[X]}(M)$ the ideal of polynomials $g(X)$ in $D[X]$ such that $g(M) = 0$, also called the null ideal of M over D.

Here is an overview of the paper. In the second section we give a characterization on the ring of polynomials which are integer-valued over the set of matrices with prescribed characteristic polynomial. If D is integrally closed, we also prove that an ideal in $D[X]$ generated by a non-zero element of D and a non-constant monic polynomial is contracted with respect to every polynomial ring extension $O_E[X]$, where O_E is the integral closure of D in a field extension of K. In the third section we recall the definition of divided difference and we use the previous results to characterize integer-valued polynomials over matrices in terms of their divided differences. In the fourth section we give some results about integer-valued polynomials over triangular matrices and the so-called polynomial closure of set of matrices having prescribed characteristic polynomial. Finally, in the last section we give some remarks.

2 Integer-valued polynomials over matrices with prescribed characteristic polynomial

Given a monic polynomial $p \in D[X]$ of degree n we consider the matrices of $M_n(D)$ whose characteristic polynomial $p_M(X)$ is equal to $p(X)$:

$$M_n^p(D) \doteqdot \{M \in M_n(D) \,|\, p_M(X) = p(X)\}.$$

We then consider the corresponding ring of integer-valued polynomials:

$$\mathrm{Int}(M_n^p(D)) \doteqdot \{f \in K[X] \,|\, f(M) \in M_n(D),\, \forall M \in M_n^p(D)\}$$

which is an overring of $\mathrm{Int}(M_n(D))$.

The following Lemma is easy to prove.

Lemma 2.1 *Let D be a commutative ring and $p \in D[X]$ a monic polynomial. Then*

$$\bigcap_{M \in M_n^p(D)} N_{D[X]}(M) = (p(X)).$$

Proof : By Cayley-Hamilton theorem we have that $p(X)$ annihilates every matrix in $M_n^p(D)$. Conversely, we use the fact that over any commutative ring D the null ideal of the companion matrix of a polynomial $p(X)$ is the principal ideal generated by $p(X)$ (see [4]). Hence, the above intersection is contained in $N_{D[X]}(C_p) = (p(X))$, so we are done. $\square$

From now on D is a domain. Given a matrix $M \in M_n(D)$, we set $\mathrm{Int}(\{M\})$ to be the ring of polynomials in $K[X]$ such that $f(M) \in M_n(D)$. It is easy to see that (see also [4]), given $f(X) = g(X)/d \in K[X]$, $g \in D[X], d \in D, d \neq 0$, we have

$$f(M) \in M_n(D) \text{ if and only if } \overline{g} \in N_{(D/dD)[X]}(\overline{M}), \tag{1}$$

where $N_{(D/dD)[X]}(\overline{M})$ is the null ideal over $(D/dD)[X]$ of the matrix $\overline{M}$ obtained by reducing the entries of M modulo the ideal dD and $\overline{g}$ is the polynomial obtained by reducing the coefficients of g modulo dD. Since $\mathrm{Int}(M_n^p(D)) = \bigcap_{M \in M_n^p(D)} \mathrm{Int}(\{M\})$ (the intersection is taken over the set of all the matrices M whose characteristic polynomial $p_M(X)$ is equal to $p(X)$), we get

$$\mathrm{Int}(M_n^p(D)) = \{f(X) = g(X)/d \in K[X] \mid \overline{g} \in \bigcap_{M \in M_n^p(D)} N_{(D/dD)[X]}(\overline{M})\}. \tag{2}$$

(notice that given $g(X)/d \in K[X]$, the reduction on the right is modulo d).

Next Lemma follows immediately from Lemma 2.1 and (2). It gives a characterization of the ring of integer-valued polynomials over the set of matrices with a prescribed characteristic polynomial.

Lemma 2.2 *Let $p \in D[X]$ be a monic polynomial of degree n and $f(X) = g(X)/d \in K[X]$. Then*

$$f \in \mathrm{Int}(M_n^p(D)) \Leftrightarrow g(X) \text{ is divisible by } p(X) \text{ modulo } dD[X].$$

Since the polynomial $p(X)$ is monic, by the Euclidean division algorithm the previous condition is equivalent to $g(X)$ belongs to the ideal of $D[X]$ generated by $p(X)$ and d. Moreover, by the aforementioned result about null ideals of companion matrices, we have $\mathrm{Int}(M_n^p(D)) = \mathrm{Int}(\{C_p\})$, where C_p is the companion matrix of $p(X)$.

We now generalize the previous definition of the set of matrices $M_n^p(D)$ with prescribed characteristic polynomial $p(X)$ to any given set S of monic polynomials in $D[X]$ of degree n. We denote by $M_n^S(D)$ the set of matrices whose characteristic polynomial is in S and by $\mathrm{Int}(M_n^S(D))$ the ring of polynomials in $K[X]$ which are integer-valued over $M_n^S(D)$, that is $\mathrm{Int}(M_n^S(D)) = \{f \in K[X] \mid f(M_n^S(D)) \subset M_n(D)\}$. Obviously, we have $M_n^S(D) = \bigcup_{p \in S} M_n^p(D)$, so that

$$\mathrm{Int}(M_n^S(D)) = \bigcap_{p \in S} \mathrm{Int}(M_n^p(D)).$$

The following Proposition follows by Lemma 2.2 and the previous representation of the ring $\mathrm{Int}(M_n^S(D))$ as an intersection of the overrings $\mathrm{Int}(M_n^p(D))$.

Proposition 2.1 *Let S be a set of monic polynomials in $D[X]$ of degree n. Let $f(X) = g(X)/d \in K[X]$. Then*

$$f \in \mathrm{Int}(M_n^S(D)) \Leftrightarrow g(X) \text{ divisible modulo } dD[X] \text{ by all } p \in S.$$

By Lemma 3.4 of [4] a polynomial $f(X) = g(X)/d \in K[X]$ is integer-valued over $M_n(D)$ if and only if $g(X)$ is divisible modulo $dD[X]$ by all monic polynomials $p \in D[X]$ of degree n. Then by Lemma 2.2 we have this representation of the ring $\mathrm{Int}(M_n(D))$:

$$\mathrm{Int}(M_n(D)) = \bigcap_{p \in \mathcal{S}_n} \mathrm{Int}(M_n^p(D)) \tag{3}$$

where $\mathcal{S}_n$ denotes the set of monic polynomials of degree n in $D[X]$. Notice that this is Proposition 2.1 for $S = \mathcal{S}_n$.

Suppose now that D has zero Jacobson radical. By Proposition 6.2 of [5] we can just check the previous condition on irreducible polynomials; namely, a polynomial $f(X) = g(X)/d \in K[X]$ is integer-valued over $M_n(D)$ if and only if $g(X)$ is divisible modulo $dD[X]$ by all monic irreducible polynomials $p \in D[X]$ of degree n. For a given integer n, we set $\mathcal{S}_n^{\mathrm{irr}}$ to be the set of monic and irreducible polynomials in $D[X]$ of degree n. We consider the set of matrices $M_n^{\mathcal{S}_n^{\mathrm{irr}}}(D) \subsetneq M_n(D)$ with irreducible characteristic polynomial. To ease the notation, we set $M_n^{\mathcal{S}_n^{\mathrm{irr}}}(D) = M_n^{\mathrm{irr}}(D)$. Then the last result shows that

$$\mathrm{Int}(M_n(D)) = \mathrm{Int}(M_n^{\mathrm{irr}}(D))$$

that is, $M_n^{\mathrm{irr}}(D)$ is polynomially dense in $M_n(D)$ (see [3] for the definition, which we recall in section 4.1). Since $M_n^{\mathrm{irr}}(D) = \bigcup_{p \in \mathcal{S}_n^{\mathrm{irr}}} M_n^p(D)$ we have this other representation of the ring $\mathrm{Int}(M_n(D))$ when the Jacobson radical of D is zero:

$$\mathrm{Int}(M_n(D)) = \bigcap_{p \in \mathcal{S}_n^{\mathrm{irr}}} \mathrm{Int}(M_n^p(D)). \tag{4}$$

2.1 A contracted ideal in $D[X]$

In this section we need the extra assumption that D is integrally closed. We make first a preliminary remark. Let E be a field extension of K and let O_E be the integral closure of D in K. Given $d \in D \setminus \{0\}$, we claim that $dO_E \cap D = dD$. In fact, if $da \in dO_E \cap D$, with $a \in O_E$, then $a = r/d$, for some $r \in D$. So a is in K, but since by assumption a is integral over D and D is integrally closed, we have that a is actually in D. The other containment is obvious. From this it also follows that $dO_E[X] \cap D[X] = dD[X]$. The containment ($\supseteq$) follows as before. For the other containment, if we have $dR(X) \in D[X]$, for some $R \in O_E[X]$, then for all the coefficients $c_i \in O_E$ of the polynomial R we have $dc_i \in D$, so that by the previous result we have $c_i \in D$. Under the assumption of D integrally closed and $p(X)$ monic, next lemma says that extending the base field does not alter the divisibility of a polynomial $g(X)$ by $p(X)$ modulo an element d of D.

Lemma 2.3 *Let D be an integrally closed domain with quotient field K. Let $p \in D[X]$ be monic and $d \in D$. Let E be a field extension of K and O_E the integral closure of D in E. Let $g \in D[X]$ be such that $g(X)$ is divisible in $O_E[X]$ by $p(X)$ modulo $dO_E[X]$. Then $g(X)$ is divisible in $D[X]$ by $p(X)$ modulo $dD[X]$.*

Equivalently, the lemma says that the ideal of $D[X]$ generated by d and $p(X)$ is a contracted ideal with respect to $O_E[X]$, that is:

$$dD[X] + p(X)D[X] = (dO_E[X] + p(X)O_E[X]) \cap D[X]$$

(notice that the inclusion ($\subseteq$) always holds).

Proof : Let n be the degree of $\deg(p)$. Let $g \in D[X]$ be such that $g \in dO_E[X] + p(X)O_E[X]$. Then we have

$$g(X) = Q(X)p(X) + dR(X)$$

for some $Q, R \in O_E[X]$. By Euclidean division ($p(X)$ is monic) we can assume that $\deg(R) < n$ (if not, just divide $R(X)$ by $p(X)$ in $O_E[X]$). We divide now $g(X)$ by $p(X)$ in $D[X]$ and we get

$$g(X) = q(X)p(X) + r(X)$$

for some $q, r \in D[X]$, $\deg(r) < n$. By the uniqueness of quotient and remainder in $O_E[X]$ we have $Q(X) = q(X)$ and $dR(X) = r(X)$. This implies that $Q \in D[X]$. By the claim before the Lemma, the second equality implies that $R \in D[X]$, and this concludes the proof. $\square$

If $d = 0$ we have $p(X)D[X] = p(X)O_E[X] \cap D[X]$. We stress that in this case we don't need the assumption that D is integrally closed, but just that $p(X)$ is monic (look at the last lines of the above proof). However, in our case d will always be non-zero, since it will be the denominator of a polynomial $f \in K[X]$, as in Lemma 2.2.

In general the statement of the Lemma is false, if we drop the assumption that D is integrally closed. For example, let $D = \mathbb{Z}[\sqrt{5}] \subset O_K = \mathbb{Z}[(1+\sqrt{5})/2] \subset K = \mathbb{Q}(\sqrt{5})$, where O_K, the ring of integers of the number field K, is the integral closure of D in its quotient field K. Take $E = K$, so that $O_E = O_K$. We have that $2D \subsetneq 2O_K \cap D = 2O_K \subset D$ (last equality holds because $2O_K$ is the conductor of the integral extension $D \subset O_K$). The element $\alpha = 1 + \sqrt{5}$ is in $2O_K$ but not in $2D$, so that the polynomial $g(X) = X + \alpha \in D[X]$ is divisible in $O_K[X]$ by X modulo $2O_K[X]$ but it is not divisible in $D[X]$ by X modulo $2D[X]$.

Under the assumptions of Lemma 2.3, given $g \in D[X]$ we have that $g(X)$ is divisible in $D[X]$ by $p(X)$ modulo $dD[X]$ if and only if $g(X)$ is divisible in $O_E[X]$ by $p(X)$ modulo $dO_E[X]$. Therefore, by extending the base field, the polynomials in $D[X]$ which are divisible in $O_E[X]$ by $p(X)$ modulo $dO_E[X]$ are only those divisible by $p(X)$ in $D[X]$ modulo $dD[X]$.

Lemma 2.2 and 2.3 show that

$$\mathrm{Int}(M_n^p(D)) = K[X] \cap \mathrm{Int}(M_n^p(O_E))$$

and we stress that the above formula holds for any field extension E of K.

We can extend the field K to a finite extension where $p(X)$ has a root or it is totally split. For example, if $p(X)$ is irreducible over K, let $E = K[X]/(p(X))$. Let F be the splitting field of $p(X)$ over K. F is equal to the compositum of the conjugates fields E_i of E over K (in particular, it depends only on E and not on the polynomial $p(X)$; the conjugate fields of E over K are just the images of E via the K-embeddings of E into $\overline{K}$). Notice that $p(X)$ splits into n linear factors over F, say $p(X) = \prod_{i=1,\ldots,n}(X - \alpha_i)$, where $\alpha_1, \ldots, \alpha_n$ are integral over D, thus they belong to O_F, the integral closure of D in F. With a suitable reordering, we may suppose that $E_i = K(\alpha_i)$, so that α_i is in the integral closure O_{E_i} of D in E_i. Notice that if the characteristic of K is not zero then $p(X)$ may have multiple roots. By Lemma 2.3 we have:

$$\mathrm{Int}(M_n^p(D)) = \mathrm{Int}(M_n^p(O_E)) \cap K[X] = \mathrm{Int}(M_n^p(O_F)) \cap K[X].$$

3 Divided differences of integer-valued polynomials over matrices

Let $g(X)$ be a polynomial of degree n over a commutative ring D and let $a_0,\dots,a_n$ be arbitrary chosen elements of D (possibly with repetitions).

We have the following formula due to Newton:

$$\begin{aligned} g(X) = g(a_0) + \Phi^1(g)(a_0,a_1)(X-a_0) + \Phi^2(g)(a_0,a_1,a_2)(X-a_0)(X-a_1) + \dots \\ + \Phi^n(g)(a_0,\dots,a_n)(X-a_0)\cdot\ldots\cdot(X-a_{n-1}) \end{aligned} \tag{5}$$

where for each $k\in\mathbb{N}$, $\Phi^k(g)(X_0,\dots,X_k)$ is the k-th divided difference of g defined recursively as follows:

$$\begin{aligned} \Phi^0(g)(X_0) &\doteqdot g(X) \\ \Phi^1(g)(X_0,X_1) &\doteqdot \frac{g(X_0)-g(X_1)}{X_0-X_1} \\ &\dots \\ \Phi^k(g)(X_0,\dots,X_k) &\doteqdot \frac{\Phi^{k-1}(g)(X_0,\dots,X_{k-1})-\Phi^{k-1}(g)(X_0,\dots,X_{k-2},X_k)}{X_{k-1}-X_k} \end{aligned}$$

$\Phi^k(g)$ is a symmetric polynomial with coefficients in D in $k+1$ variables. Obviously, a polynomial $g\in D[X]$ is divisible by $(X-a_0)\cdot\ldots\cdot(X-a_{n-1})$ if and only if the coefficients $\Phi^k(g)(a_0,\dots,a_k)$ of the expansion in (5) are equal to zero, for all $0\le k<n$. We just use the fact that $\{1\}\cup\{\prod_{i=0,\dots,k}(X-a_i)\}_{k=0,\dots,n-1}$ are linearly independent over D. More in general, if $\{a_i\}_{i\in\mathbb{N}}$ is a sequence in D, then $\{1\}\cup\{\prod_{i=0,\dots,k}(X-a_i)\}_{k\in\mathbb{N}}$ is a free basis of the D-module $D[X]$.

Next lemma is a generalization of the previous remark and gives the following criterion: a polynomial $g(X)$ is divisible by a totally split polynomial $p(X)$ modulo a principal ideal $dD[X]$ if and only if the evaluation of the divided differences of $g(X)$ at the roots of $p(X)$ are divisible by d. Notice that by Lemma 2.2, the polynomial $g(X)/d$ is in $\mathrm{Int}(M_n^p(D))$. We will use this criterion together with Lemma 2.3, which allows us to extend the base field where the polynomial $p(X)$ is totally split. This will lead to the characterization about the divided differences of an integer-valued polynomial over $M_n(D)$ we talked about in the introduction. Next lemma appears also in [3], but for the sake of the reader we report it here.

Lemma 3.1 *Let D be a domain. Let $g\in D[X]$, $d\in D\setminus\{0\}$ and $a_0,\dots,a_n\in D$ (not necessarily distinct). Then $g(X)$ is divisible modulo $dD[X]$ by $p(X)=\prod_{i=0,\dots,n}(X-a_i)$ if and only if for all $0\le k<n$ we have $\Phi^k(g)(a_0,\dots,a_k)\in dD$.*

Proof : Consider everything modulo dD and apply Newton's formula. $\square$

Let D be an integrally closed domain with quotient field K. Remember that for any extension E of K we denote by O_E the integral closure of D in E. From now on we consider only algebraic extension of K, which are tacitly assumed to be contained in $\overline{K}$. We give now the following proposition:

Proposition 3.1 *Let D be an integrally closed domain with quotient field K. Let $g \in D[X]$, $d \in D \setminus \{0\}$ and $p \in D[X]$ monic of degree n. Let $A_p = \{\alpha_1, \dots, \alpha_n\}$ be the set of roots of $p(X)$ in a splitting field F over K. Then $f(X) = g(X)/d \in \mathrm{Int}(M_n^p(D))$ if and only if for all $0 \leq k < n$, $\Phi^k(g)(\alpha_{i_0}, \dots, \alpha_{i_k}) \in dO_F$, for all subsets $\{i_0, \dots, i_k\}$ of $k+1$ elements of $\{1, \dots, n\}$.*

Notice that when we write $\{\alpha_1, \dots, \alpha_n\}$ this is a multi-set, in the sense we allow repetions, in the case of multiple roots of the polynomial $p(X)$. In the above statement we can choose a root α_i of $p(X)$ multiple times if it has multiplicity greater than 1 (we are not allowed to choose a root more times than its multiplicity, though). So indeed $\{a_{i_0}, \dots, a_{i_k}\}$ can be a multi-set.

Proof : By Lemma 2.2, $f(X) = g(X)/d$ is in $\mathrm{Int}(M_n^p(D))$ if and only if $g(X)$ is divisible modulo $dD[X]$ by $p(X)$. Since D is integrally closed, by Lemma 2.3 this is equivalent to $g(X)$ divisible modulo $dO_F[X]$ by $p(X)$. If we fix an order of the roots $\alpha_1, \dots, \alpha_n$ of $p(X)$, by Newton's formula (5) and Lemma 3.1, the previous condition holds if and only if $\Phi^k(g)(\alpha_0, \dots, \alpha_k) \in dO_F$ for all $0 \leq k < n$. Since this holds for every possible order of the α_i's we may choose (essentially because $\Phi^k(g)(X_1, \dots, X_{k+1})$ is symmetric), we get the final statement. $\square$

Notice that the roots of $p(X)$ are integral over D (hence they are in O_F) and by definition $F = K(\alpha_1, \dots, \alpha_n)$. We stress that we are not assuming $p(X)$ to be irreducible. We give now the following definition (remember that $\overline{D}$ is the integral closure of D in $\overline{K}$):

Definition: Let $f \in K[X_1, \dots, X_n]$. Let $\alpha_1, \dots, \alpha_n \in \overline{D}$. We say that $f(\underline{X})$ is integral on the n-tuple $(\alpha_1, \dots, \alpha_n) \in \overline{D}^n$ if $f(\alpha_1, \dots, \alpha_n)$ is integral over D. We say that $f(\underline{X})$ is integral on the multi-set $\{\alpha_1, \dots, \alpha_n\} \subset \overline{D}$ if $f(\underline{X})$ is integral on $(\alpha_{i_1}, \dots, \alpha_{i_n})$, for every possible order of the α_i's we may choose.

We stress that in the previous definition we are neither assuming $\alpha_1, \dots, \alpha_n$ to be distinct nor to be conjugate over K. If the polynomial $f(\underline{X})$ is symmetric on its variables (like the divided differences) then, if $f(\underline{X})$ is integral on $(\alpha_1, \dots, \alpha_n)$, then $f(\underline{X})$ is integral on $\{\alpha_1, \dots, \alpha_n\}$. If $f(\alpha_1, \dots, \alpha_n)$ is integral over D, then it obviously belongs to the integral closure of D in the field $K(\alpha_1, \dots, \alpha_n)$ generated by $\alpha_1, \dots, \alpha_n$ over K inside $\overline{K}$.

Using these definitions, the statement of Proposition 3.1 is equivalent to the following. For all $0 \leq k < n$, $\Phi^k(f)$ is integral on every subset $\{\alpha_{i_0}, \dots, \alpha_{i_k}\}$ of the roots A_p of $p(X)$. In particular, for each of the relevant k, $\Phi^k(f)(\alpha_{i_0}, \dots, \alpha_{i_k}) \in O_{F_k}$, where F_k is the field generated by $\alpha_{i_0}, \dots, \alpha_{i_k}$ over K. By the argument given in the proof of Proposition 3.1, if we fix an ordering of the roots A_p of $p(X)$, for $f(X)$ to belong to $\mathrm{Int}(M_n^p(D))$ it is sufficient that each k-th divided difference of $f(X)$ is integral on the first $k+1$ roots of $p(X)$.

Let us see with an example what happens in the case $n = 2$. Let $p \in D[X]$ be an irreducible monic polynomial of degree 2. Let α be a root of $p(X)$ in $\overline{K}$. Over the quadratic extension $K(\alpha)$ of K we have $p(X) = (X - \alpha)(X - \alpha')$, where α' is the conjugate root of α. Let $O_{K(\alpha)}$ be the integral closure of D in $K(\alpha)$. Then, given a polynomial $f(X)$ in $K[X]$ we have that $f(X)$ is integer-valued over the set of matrices in $M_2(D)$ with characteristic polynomial equal to $p(X)$ if and only if $f(\alpha), f(\alpha') \in O_{K(\alpha)}$ and $\Phi^1(f)(\alpha, \alpha') \in O_{K(\alpha)}$. If the extension $K(\alpha) \supset K$ is separable, then $\Phi^1(f)(\alpha, \alpha') \in D$, because $\Phi^1(f)(X, Y)$ is symmetric, so in particular $\Phi^1(f)(\alpha, \alpha')$ is invariant under the action of the Galois group of $K(\alpha)$ over K. If we consider a split polynomial $p(X)$ in

$D[X]$, say $p(X) = (X-n)(X-m)$, for some $n, m \in D$, then $f \in \mathrm{Int}(M_2^p(D))$ if and only if $f(n), f(m) \in D$ and $\Phi^1(f)(n,m) \in D$.

Finally, by Proposition 3.1 and the representation of $\mathrm{Int}(M_n(D))$ in (3) and (4), we give this characterization of the polynomials of $\mathrm{Int}(M_n(D))$ in terms of their divided differences. If the integrally closed domain D has also zero Jacobson radical, we obtain a less restrictive condition.

Theorem 3.1 *Let D be an integrally closed domain. Let $f \in K[X]$ and n a positive integer. Then $f \in \mathrm{Int}(M_n(D))$ if and only if for every $0 \leq k < n$, for every monic polynomial $p \in D[X]$ of degree n and for every set of $k+1$ elements $\{\alpha_1, \ldots, \alpha_{k+1}\}$ of the roots of $p(X)$ in $\overline{K}$, $\Phi^k(f)$ is integral on $\{\alpha_1, \ldots, \alpha_{k+1}\}$.*

If D has also zero Jacobson radical then it is sufficient to consider polynomials $p(X)$ which are irreducible.

The roots α_i of $p(X)$ have degree bounded by $n = \deg(p)$. Notice that the divided differences of an integer-valued polynomial over $M_n(D)$ are integral on subsets $\{\alpha_1, \ldots, \alpha_{k+1}\} \subset \overline{D}$ of the roots of monic polynomials of degree n with coefficients in D, that is, the integral elements $\{\alpha_1, \ldots, \alpha_{k+1}\}$ of degree $\leq n$ cannot be chosen independently (except when they lie in D, see final remarks in the last section; this happens if $p(X)$ is totally split over D).

4 Integer-valued polynomials over triangular matrices with prescribed characteristic polynomial

We give here some characterizations of integer-valued polynomials over triangular matrices. Now D is again just an integral domain. Given a monic totally split polynomial $p \in D[X]$ (that is, $p(X) = \prod_{i=1,\ldots,n}(X - a_i)$, with $a_i \in D$), we denote by $T_n^p(D)$ the set of triangular matrices with characteristic polynomial equal to $p(X)$. This means that the entries in the diagonal of T are the roots of $p(X)$. We denote by $\mathrm{Int}(T_n^p(D))$ the ring of polynomials in $K[X]$ which are integer-valued over $T_n^p(D)$, that is $\mathrm{Int}(T_n^p(D)) = \{f \in K[X] \mid f(T_n^p(D)) \subset M_n(D)\}$ (notice that the image of a triangular matrix under a polynomial is still a triangular matrix). Next lemma is analogous to Proposition 14 of [3].

Lemma 4.1 *Let $p \in D[X]$ be a totally split polynomial of degree n. Let $f(X) = g(X)/d$ be in $K[X]$. Then*

$$f \in \mathrm{Int}(T_n^p(D)) \Leftrightarrow g(X) \text{ is divisible by } p(X) \text{ modulo } dD[X].$$

Proof : One direction is easy (the proof is exactly like in Lemma 2.2): suppose $g(X) = h(X)p(X) + dr(X)$, for some $h, r \in D[X]$. Let T be a triangular matrix with characteristic polynomial equal to $p(X)$. Then by the Cayley-Hamilton theorem, we have that $g(T) = dr(T)$, so that $f(T) = r(T) \in M_n(D)$.

For the converse, we use an inductive argument as in [3], Proposition 14. The induction is on the degree n of $p(X)$. Let $n = 2$ and let $p(X) = (X - a_0)(X - a_1)$, for some $a_0, a_1 \in D$. Given $f(X) = g(X)/d$ in $\mathrm{Int}(T_2^p(D))$, we divide $g(X)$ by $p(X)$ in $D[X]$: $g(X) = q(X)p(X) + r(X)$, with

$q, r \in D[X]$, $\deg(r) \le 1$, say $r(X) = aX + b$. Take

$$T = \begin{pmatrix} a_0 & 0 \\ 1 & a_1 \end{pmatrix}.$$

Then $g(T) = aT + b \in T_2(dD)$ (essentially by (1)) is the triangular matrix

$$\begin{pmatrix} aa_0 + b & 0 \\ a & aa_1 + b \end{pmatrix}$$

so that $a, b \in dD$ as wanted.

Suppose now the statement is true up to n and suppose that $p(X) = \prod_{i=0,\dots,n}(X - a_i)$. Let $p_1(X) = \prod_{i=1,\dots,n}(X - a_i)$. Given $T \in T_n^{p_1}(D)$, we consider the following $(n+1) \times (n+1)$ matrix:

$$T_1 = \left(\begin{array}{c|c} \text{T} & \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \\ \hline 0 \cdots 0 & a_0 \end{array} \right).$$

Notice that T_1 is in $T_{n+1}^p(D)$ and it is a block matrix. Let $f \in \operatorname{Int}(T_{n+1}^p(D))$. Then $f(X)$ is integer-valued on T. In fact,

$$f(T_1) = \left(\begin{array}{c|c} \text{f(T)} & \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} \\ \hline 0 \cdots 0 & f(a_0) \end{array} \right)$$

$f(T_1)$ has entries in D, so that $f(T)$ has entries in D, too. Since this holds for every T in $T_n^{p_1}(D)$ and for every $f \in \operatorname{Int}(T_{n+1}^p(D))$, we have $\operatorname{Int}(T_{n+1}^p(D)) \subset \operatorname{Int}(T_n^{p_1}(D))$.

By induction hypothesis, given $f(X) = g(X)/d \in \operatorname{Int}(T_{n+1}^p(D))$, we have that $g(X)$ is divisible by $p_1(X)$ modulo $dD[X]$. Then we continue as in the proof of Proposition 14 of [3], essentially going through the characterization of Lemma 3.1 together with Corollary 13 of [3]. □

In particular, Lemma 2.2 and 4.1 show that

$$\operatorname{Int}(T_n^p(D)) = \operatorname{Int}(M_n^p(D)).$$

Given a set S of monic totally split polynomials in $D[X]$ of degree n, we denote by $T_n^S(D)$ the set of triangular matrices with characteristic polynomial in S. The following proposition follows from the obvious fact $T_n^S(D) = \bigcup_{p \in S} T_n^p(D)$. This is the analogous of Proposition 2.1.

Proposition 4.1 *Let S be a set of monic totally split polynomials in $D[X]$ of degree n. Let $f(X) = g(X)/d \in K[X]$. Then*

$$f \in \operatorname{Int}(T_n^S(D)) \Leftrightarrow g(X) \text{ divisible modulo } dD[X] \text{ by all } p \in S.$$

In particular,

$$\operatorname{Int}(T_n^S(D)) = \operatorname{Int}(M_n^S(D)).$$

Last statement follows from Proposition 2.1.

4.1 Polynomial closure of set of matrices

Given a subset $\mathcal{M}$ in $M_n(D)$, the polynomial closure of $\mathcal{M}$ in $M_n(D)$ (as defined in [3]) is the set

$$\overline{\mathcal{M}} \doteqdot \{M \in M_n(D) \mid f(M) \in M_n(D),\ \forall f \in \mathrm{Int}(\mathcal{M})\}$$

where $\mathrm{Int}(\mathcal{M}) = \{f \in K[X] \,|\, f(\mathcal{M}) \subset M_n(D)\}$. $\overline{\mathcal{M}}$ is the largest set in $M_n(D)$ containing $\mathcal{M}$ such that $\mathrm{Int}(\mathcal{M}) = \mathrm{Int}(\overline{\mathcal{M}})$. A set of matrices $\mathcal{M}$ is said to be polynomially closed if $\mathcal{M} = \overline{\mathcal{M}}$.

For some domains D, the authors of [3] prove that, if S is equal to the set $\mathcal{S}_n^s$ of all monic totally split polynomials in $D[X]$ of degree n, the polynomial closure of $T_n^{\mathcal{S}_n^s}(D) = T_n(D)$ is equal to the set $M_n^{\mathcal{S}_n^s}(D)$ of all matrices in $M_n(D)$ with totally split characteristic polynomial (see Theorem 24 in [3], where the above set of matrices is denoted by $S_n(D)$). On the other hand, Proposition 4.1 above shows that, for any given subset $S \subset \mathcal{S}_n^s$, $T_n^S(D)$ and $M_n^S(D)$ have the same polynomial closure in $M_n(D)$, where D is any integral domain. So we may restrict our attention to the polynomial closure of set of matrices of the form $M_n^S(D)$, rather than just a set $T_n^S(D)$ of triangular matrices.

Under a rather general assumption, we characterize the polynomial closure $\overline{M_n^p(D)}$ of $M_n^p(D)$ in $M_n(D)$, for any given monic polynomial $p \in D[X]$ of degree n. If $p(X)$ is totally split over D, then $\overline{M_n^p(D)} = \overline{T_n^p(D)}$, by what we have just said. We make first a remark. Given any non-zero element $d \in D$, we set $\mathcal{M}_d$ to be the set of maximal ideals of D containing d. If d is a unit then $\mathcal{M}_d$ is the empty set. It is clear that the maximal spectrum of D is equal to $\bigcup_{d \in D\setminus\{0\}} \mathcal{M}_d$ (there is no maximal ideal containing only 0, except in the trivial case that D is a field, which we are clearly excluding in our considerations). From this it follows that

$$\bigcap_{d \in D\setminus\{0\}} dD \subset \bigcap_{\mathfrak{M} \subset D} \mathfrak{M} \tag{6}$$

where the latter intersection is taken over the set of all maximal ideals $\mathfrak{M}$ of D, that is, it is the Jacobson radical of D. We denote by $\mathcal{P}$ the intersection of all the non-zero principal ideals of D, that is $\bigcap_{d \in D\setminus\{0\}} dD$.

Lemma 4.2 *Suppose D is an integral domain with zero Jacobson radical. Let $p \in \mathcal{S}_n$. Then the polynomial closure of $M_n^p(D)$ is equal to the set of matrices annihilated by $p(X)$.*

Proof : By Lemma 2.2, a polynomial $f(X)$ in $\mathrm{Int}(M_n^p(D))$ has the form $f(X) = r(X) + p(X)g(X)/d$, for some $r, g \in D[X]$, $d \in D \setminus \{0\}$ (since $p(X)$ is monic, we can also assume $\deg(r) < \deg(p)$).

Clearly, if $p(M) = 0$ then for every polynomial $f \in \mathrm{Int}(M_n^p(D))$ as above, we have $f(M) = r(M) \in M_n(D)$. Conversely, a matrix M is in the polynomial closure of $M_n^p(D)$ if and only if, for every $d \in D \setminus \{0\}$ and for every $g \in D[X]$, we have $p(M)g(M) \in M_n(dD)$. In particular, this implies that $p(M) \in M_n(\mathcal{P})$. Since D has zero Jacobson radical, by (6) we have that $p(M) = 0$ as wanted. □

Without the above assumption on D, the polynomial closure of $M_n^p(D)$ is equal to the set $\{M \in M_n(D) \,|\, p(M) \in M_n(\mathcal{P})\}$.

For a polynomial $p \in \mathcal{S}_n$, like before we denote by A_p the set of roots of $p(X)$ in the algebraic closure $\overline{K}$ of the quotient field K of D. Notice that $A_p \subset \overline{D}$, the integral closure of D in $\overline{K}$.

Corollary 4.1 *Let D be an integral domain with zero Jacobson radical. Let $p \in \mathcal{S}_n$ and let $M \in M_n(D)$ such that $A_{p_M} \not\subset A_p$. Then M is not contained in the polynomial closure of $M_n^p(D)$.*

Proof : Suppose by contradiction that M is contained in the polynomial closure of $M_n^p(D)$. By previous Lemma we have $p \in N_{D[X]}(M)$. By a theorem of McCoy (see Corollary pg.283 of [6]) the ideals $N_{D[X]}(M)$ and $(p_M(X))$ of $D[X]$ have the same radical, so that $p(X)^m \in (p_M(X))$ for some $m \in \mathbb{N}$. Then for all the roots α of $p_M(X)$ we have $p(\alpha) = 0$ which is not possible by assumption. $\square$

Corollary 4.2 *Let D be an integral domain with zero Jacobson radical. Let $p \in \mathcal{S}_n$ be such that $p(X)$ is irreducible in $K[X]$. Then $M_n^p(D)$ is polynomially closed.*

If Gauss's lemma applies, we may just suppose that $p(X)$ is irreducible in $D[X]$.

Proof : By Lemma 4.2, $\overline{M_n^p(D)} = \{M \in M_n(D) \,|\, p(M) = 0\}$. Let $M \in M_n(D)$ be a matrix annihilated by $p(X)$. Since $p(X)$ is irreducible it is equal to the minimal polynomial of M in $K[X]$, that is $p(X)$ is the generator of $N_{K[X]}(M)$. Then the characteristic polynomial of M (considered over D or over K is the same, since M has entries in D) is equal to the minimal polynomial of M, because they have the same irreducible factors. Then $p(X)$ is the characteristic polynomial of M, that is, M is in $M_n^p(D)$. $\square$

We leave the following open problem.
Problem: given any subset S in $\mathcal{S}_n$, compute $\overline{M_n^S(D)}$.

Since $M_n^S(D) = \bigcup_{p \in S} M_n^p(D)$, by Lemma 4.2 and known properties of polynomial closure (see [2]) we have that

$$\bigcup_{p \in S} \overline{M_n^p(D)} = \{M \in M_n(D) \,|\, p(M) = 0, \text{ for some } p \in S\} \subset \overline{M_n^S(D)}.$$

It is not true in general that the above containment is an equality. For example, let $D = \mathbb{Z}$ and let K be a quadratic number field with ring of integers O_K. Let S_K be the set of minimal polynomials of all the quadratic algebraic integers in O_K and let $M_2^K(\mathbb{Z})$ be the set of matrices $M_2^{S_K}(\mathbb{Z})$. Then by Theorem 4.1 of [7] we have that $\mathrm{Int}(M_2^K(\mathbb{Z})) \subset \mathrm{Int}_{\mathbb{Q}}(O_K) \subset \mathrm{Int}(\mathbb{Z})$. In particular this implies that any 2×2 diagonal matrix D with rational integers in the diagonal is contained in the polynomial closure of $M_2^K(\mathbb{Z})$. Obviously the null ideal of such a matrix D over $\mathbb{Z}$ is generated by a totally split polynomial in $\mathbb{Z}$, which does not belong to S_K.

5 Final Remarks

Let $f(X)$ be in $\mathrm{Int}(M_n(D))$. By Theorem 3.1, for each $0 \leq k \leq n-1$, the k-th divided difference $\Phi^k(f)$ is integral on every set $\{\alpha_0, \ldots, \alpha_k\}$ of roots of any monic polynomial $p(X)$ in $D[X]$ of degree n (these roots lies in the algebraic closure $\overline{K}$ of the quotient field K of D; there may be repetitions among these roots). In particular, $\Phi^{n-1}(f)$ is integral-valued on every full set of conjugate integral elements of degree n, that is, the roots of an irreducible monic polynomial $p(X)$ of degree n. If

$p(X)$ is separable, the corresponding value of $\Phi^{n-1}(f)$ on the roots of $p(X)$ lies in D, by Galois invariance (the divided differences of $f(X)$ all have coefficients in K).

If we consider polynomials which split into linear factors in $D[X]$ we see that $\Phi^k(f)$ is integer-valued on every multi-set $\{\alpha_0, \dots, \alpha_k\}$ of elements of D. Thus over D the elements $\alpha_0, \dots, \alpha_k$ can be chosen independently. Hence, $\Phi^k(f)(D^{k+1}) \subset D$. This implies that $\mathrm{Int}(M_n(D))$ is a subring of the ring of integer-valued polynomials whose k-th divided differences are also integer-valued for all $k < n$. This ring has been introduced by Bharghava in [1] and it is denoted by:

$$\mathrm{Int}^{\{n-1\}}(D) \doteqdot \{f \in K[X] \,|\, \Phi^k(f)(D^{k+1}) \subset D\,, \forall 0 \leq k < n\}.$$

As we said in the introduction, in [3] the authors prove that the latter ring is equal to the ring of polynomials which are integer-valued over the set of triangular matrices over D.

Moreover, for every $0 \leq k < n$, $\Phi^k(f)$ is integral-valued on every full set of roots of any monic irreducible polynomial $q(X)$ in $D[X]$ of degree $k+1$ (that is, a set of $k+1$ conjugate integral elements of degree $k+1$ over D; given such a polynomial $q(X)$ multiply it by $X^{n-(k+1)}$). We set

$$\widetilde{A_k^k} \doteqdot \{(\alpha_1, \dots, \alpha_k) \in A_k^k \,|\, \alpha_i\text{'s are conjugates}\}$$

where A_k denotes the set of integral elements over D of degree k in $\overline{K}$. Then $\Phi^{k-1}(f)(\widetilde{A_k^k})$ is contained in the set $\mathcal{A}_k$ of all the integral elements in $\overline{K}$ of degree k over K.

For $k = 0$ we see that $f(X)$ is integral on any integral element α of degree $\leq n$, so that $f \in \mathrm{Int}_K(O_E) \doteqdot K[X] \cap \mathrm{Int}(O_E)$, for every finite extension E of K of degree $\leq n$. $\mathrm{Int}(O_E)$ is the subring of $E[X]$ of those polynomials $f(X)$ such that $f(O_E) \subset O_E$. In this way we have:

$$\mathrm{Int}(M_n(D)) \subset \bigcap_{[E:K] \leq n} \mathrm{Int}_K(O_E).$$

We can easily prove that the latter intersection is equal to $\mathrm{Int}_K(\mathcal{A}_n) \doteqdot \{f \in K[X] \,|\, f(\mathcal{A}_n) \subset \mathcal{A}_n\}$, where $\mathcal{A}_n$ is the set of all the integral elements over D in $\overline{D}$ of degree bounded by n (see also [7], in the case $D = \mathbb{Z}$).

Finally, if D has also zero Jacobson radical, in order for a polynomial $f \in K[X]$ to be integer-valued over $M_n(D)$ it is sufficient to check that, for each $0 \leq k \leq n-1$, the k-th divided difference of $f(X)$ is integral on every set of $k+1$ conjugate elements of degree n over D inside $\overline{K}$ (that is, the roots of a monic irreducible polynomial of degree n in $D[X]$).

Acknowledgments

I wish to thank Keith Johnson for the useful suggestions. The author was supported by the Austrian Science Foundation (FWF), Project Number P23245-N18.

References

[1] M. Bhargava, *On P-orderings, Rings of Integer-Valued Polynomials and Ultrametric Analysis*, Journal of the Amer. Math. Soc. 22 (4) (2009), 963-993.

[2] J.-P. Cahen, J.-L. Chabert, *Integer-valued polynomials*, Amer. Math. Soc. Surveys and Monographs, 48, Providence, 1997.

[3] S. Evrard, Y. Fares, K. Johnson. *Integer valued polynomials on lower triangular integer matrices*, to appear in Monats. für Math..

[4] S. Frisch. *Polynomial Separation of Points in Algebras*, S. Chapman (ed.), Arithmetical Properties of Commutative Rings and Modules (Chapel Hill Conf.), Dekker 2005, pp. 249-254.

[5] S. Frisch. *Integer-valued polynomials on algebras*, J. of Algebra 373 (2013), 414-425.

[6] N. H. McCoy. *Concerning matrices with elements in a commutative ring*, Bull. Amer. Math. Soc. 45 (1939), no.4, 280-284.

[7] G. Peruginelli. *Integral-valued polynomials over sets of algebraic integers of bounded degree*, submitted, http://arxiv.org/abs/1301.2045.